

Relationship between ϖ and redshift z

In principle

- R-W metric:

$$d_p(t) = R(t) \int_0^{\varpi} \frac{d\varpi'}{\sqrt{1 - k\varpi'^2}}$$

Proper Distance

$$\int_e^t \frac{cdt}{R(t)} = \int_0^{\tilde{\omega}} \frac{d\tilde{\omega}}{\sqrt{1 - k\tilde{\omega}^2}}$$

$$\int_{R(t_e)}^{R(t)} \frac{c dR}{R \frac{dR}{dt}} = \int_0^{\tilde{\omega}} \frac{d\tilde{\omega}}{\sqrt{1 - k\tilde{\omega}^2}}$$

- Friedmann Eqn:

$$\rho(t) = \rho_m(t) + \rho_{rel}(t) + \rho_\Lambda(t)$$

$$\left(\frac{dR}{dt}\right)^2 - \frac{8}{3}\pi G\rho(t)R^2(t) = -kc^2$$

$$\frac{dR}{dt} = \sqrt{\frac{8}{3}\pi G\rho(t)R^2(t) - kc^2}$$

also: $R(t_e) = 1 \quad R(t) = \frac{1}{1+z}$

$$\int_{\frac{1}{1+z}}^1 \frac{c dR}{R \dots} = \int_0^{\tilde{\omega}} \frac{d\tilde{\omega}}{\sqrt{1 - k\tilde{\omega}^2}}$$

In practice

(because of that @#\$% cosmological constant)

$$I(z) = H_0 \int_{z_0}^1 \frac{dR}{R(dR/dt)} = H_0 \int_0^z \frac{dz'}{H(z')}. \quad (29.167)$$

Using Eq. (29.122), we obtain

$$I(z) = \int_0^z \frac{dz'}{\sqrt{\Omega_{m,0}(1+z')^3 + \Omega_{rel,0}(1+z')^2 + \Omega_{\Lambda,0} + (1-\Omega_0)(1+z')^2}}. \quad (29.168)$$

With this definition of the integral $I(z)$, the present proper distance is

$$d_{p,0}(z) = \frac{c}{H_0} I(z). \quad (29.169)$$

$$\varpi(z) = \frac{c}{H_0} S(z)$$

$$S(z) = I(z) \quad (\Omega_0 = 1)$$

$$= \frac{1}{\sqrt{\Omega_0 - 1}} \sin \left[I(z) \sqrt{\Omega_0 - 1} \right] \quad (\Omega_0 > 1)$$

$$= \frac{1}{\sqrt{1 - \Omega_0}} \sinh \left[I(z) \sqrt{1 - \Omega_0} \right] \quad (\Omega_0 < 1).$$


$$I(z) = \int_0^z \left\{ 1 - (1+q_0)z' + \left[\frac{1}{2} + 2q_0 + \frac{3}{2}q_0^2 + \frac{1}{2}(1-\Omega_0) \right] z'^2 + \dots \right\} dz' \quad (29.178)$$

$$I(z) = z - \frac{1}{2}(1+q_0)z^2 + \left[\frac{1}{6} + \frac{2}{3}q_0 + \frac{1}{2}q_0^2 + \frac{1}{6}(1-\Omega_0) \right] z^3 + \dots \quad (29.179)$$

$$d_{p,0} \sim \varpi \simeq \frac{cz}{H_0} \left[1 - \frac{1}{2}(1+q_0)z \right] \quad (\text{for } z \ll 1).$$

Eqns. 29.180, 29.181, true for all models

Luminosity Distance



$$F = \frac{L}{4\pi d^2}$$

$$F = \frac{L}{4\pi \varpi^2 (1+z)^2}$$

Redshift $\rightarrow (1+z)$
Time dilation $\rightarrow (1+z)$

$$d_L = \varpi (1+z)$$

About right...

$$d_L(z) \simeq \frac{cz}{H_0} \left[1 + \frac{1}{2}(1 - q_0)z \right] \quad (\text{for } z \ll 1)$$

$$m - M = 5 \log_{10}(d_L/10 \text{ pc})$$

$$m - M \simeq 5 \log_{10} \left[\frac{c}{(100 \text{ km s}^{-1} \text{ Mpc}^{-1})(10 \text{ pc})} \right] - 5 \log_{10}(h) + 5 \log_{10}(z) + 5 \log_{10} \left[1 + \frac{1}{2}(1 - q_0)z \right] \quad (\text{for } z \ll 1).$$

$$m - M = 5 \log_{10} \left[\frac{c}{(100 \text{ km s}^{-1} \text{ Mpc}^{-1})(10 \text{ pc})} \right] - 5 \log_{10}(h) + 5 \log_{10}(1+z) + 5 \log_{10}[S(z)] = 42.38 - 5 \log_{10}(h) + 5 \log_{10}(1+z) + 5 \log_{10}[S(z)].$$

$$m - M \simeq 42.38 - 5 \log_{10}(h) + 5 \log_{10}(z) + 1.086(1 - q_0)z \quad (\text{for } z \ll 1). \quad (29.188)$$

In practice

(because of that @#\$% cosmological constant)


$$d_L(z) = \frac{c}{H_0} (1+z) S(z)$$

$$S(z) = I(z) \quad (\Omega_0 = 1)$$

$$= \frac{1}{\sqrt{\Omega_0 - 1}} \sin \left[I(z) \sqrt{\Omega_0 - 1} \right] \quad (\Omega_0 > 1)$$


$$= \frac{1}{\sqrt{1 - \Omega_0}} \sinh \left[I(z) \sqrt{1 - \Omega_0} \right] \quad (\Omega_0 < 1).$$

Old Redshift-Magnitude Results



q_0 – the accelerating universe

- Type Ia supernovae are best standard candles.
 - Least scatter in luminosity
- 2 independent groups get same answer
 - Supernova Cosmology Project
 - Perlmutter et al. 1999 ApJ 517, 565
 - High- z Supernova Search
 - Garnavich et al. ApJ 509, 74
- Found *acceleration*
 - Not deceleration as expected.



[CO fig. 29.27]

